



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

For the center of gravity of the *curved* surface we have,

$$\bar{x} = \frac{4 \int xy ds}{S}, = \frac{8r \int_0^r x dx}{8r^2} = \frac{1}{r} \int_0^r x dx, = \frac{1}{2}r, = \frac{1}{2}a.$$

For the center of gravity of the *whole* surface, since the curved surface is *twice* that of the base we have, $\bar{x} = \frac{2}{3} \cdot \frac{1}{2}a = \frac{1}{3}a$.

Also solved by H. C. WHITAKER, C. W. M. BLACK, and the PROPOSER.

Professors Black and Scheffer used "side= $2a$ " as in Problem 47, instead of *side*= a , and hence their results did not agree with those in the published solutions. The results obtained were : Volume= $8a^3/3$, surface= $8a^2$, center of gravity of volume= $3a/8$, and center of gravity of surface= $\frac{1}{2}a$. See problem 42 for two additional SOLUTIONS for surface and volume. EDITOR.

MECHANICS.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

38. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.

A prolate spheroid of revolution is fixed at its focus ; a blow is given it at the extremity of the axis minor in a line tangent to the direction perpendicular to the axis major. Find the axis about which the body begins to rotate. [From *Loudon's Rigid Dynamics*.]

Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas-Texas.

The general equations of motion are :

$$\left. \begin{aligned} A\omega_x - (\Sigma mxy)\omega_y - (\Sigma mxz)\omega_z &= L \\ B\omega_y - (\Sigma myz)\omega_z - (\Sigma myx)\omega_x &= M \\ C\omega_z - (\Sigma mzx)\omega_x - (\Sigma mzy)\omega_y &= N \end{aligned} \right\} \dots\dots\dots (1).$$

The equation to the ellipsoid with focus as origin is $a^2y^2 + a^2z^2 + b^2x^2 = 2aeb^2x + b^4$. Now $\Sigma mxy = \Sigma mxz = \Sigma myz = 0$. \therefore (1) reduce to

$$\left. \begin{aligned} A\omega_x &= L \\ B\omega_y &= M \\ C\omega_z &= N \end{aligned} \right\} \dots\dots\dots (2).$$

Let $2aeb^2x + b^4 - b^2x^2 = a^2c^2$. Then

$$\begin{aligned}
 A = \Sigma m(y^2 + z^2) &= \mu \int_{-a(1-e)}^{a(1+e)} \int_{-c}^c \int_{-\sqrt{c^2-y^2}}^{\sqrt{c^2-y^2}} (y^2 + z^2) dx dy dz \\
 &= \frac{4\mu}{3} \int_{-a(1-e)}^{a(1+e)} \int_0^c \{3y^2 \sqrt{c^2-y^2} + (c^2-y^2) \sqrt{c^2-y^2}\} dx dy \\
 &= \frac{\pi\mu}{2a^4} \int_{-a(1-e)}^{a(1+e)} (2aeb^2x + b^4 - b^2x^2)^2 dx = \frac{8}{15} \mu \pi a b^4.
 \end{aligned}$$

$$\begin{aligned}
 B = C = \Sigma m(x^2 + y^2) &= \mu \int_{-a(1-e)}^{a(1+e)} \int_{-c}^c \int_{-\sqrt{c^2-y^2}}^{\sqrt{c^2-y^2}} (x^2 + y^2) dx dy dz \\
 &= 4\mu \int_{-a(1-e)}^{a(1+e)} \int_0^c (x^2 + y^2) \sqrt{c^2-y^2} dx dy \\
 &= \frac{\mu\pi}{4} \int_{-a(1-e)}^{a(1+e)} (4c^2x^2 + c^4) dx = \frac{8}{15} \mu \pi a^3 b^2 (1 + 2e^2).
 \end{aligned}$$

Let the blow $= P$ be struck perpendicular to the plane (xy) , then the moments of the impulsive forces about the axes are $L = Pb$, $M = Pa e$, $N = 0$.

These in (2) give

$$\left. \begin{aligned} \frac{8}{15} \mu \pi a b^4 \omega_x &= Pb \\ \frac{8}{15} \mu \pi a^3 b^2 (1 + 2e^2) \omega_y &= Pa e \\ \omega_z &= 0 \end{aligned} \right\} \dots\dots\dots (3).$$

$$\therefore \frac{\omega_y}{\omega_x} = \frac{e^2}{1 + 2e^2} \cdot \frac{b}{ae}.$$

Let F be the focus, O the center of the ellipsoid. Then on the minor axis in the plane (xy) , take $OE = \frac{e^2}{1 + 2e^2} \cdot b$, then will FE be the axis required.

Let $a = 5$, $b = 4$. $\therefore e = \frac{3}{5}$. $\therefore OE = \frac{9}{43} b = \frac{36}{43}$. The resultant angular velocity will be

$$\frac{15P}{8\mu\pi a^2 b^3 e} \cdot OF = \frac{P}{512\mu\pi} \cdot OF = \frac{3P\sqrt{1993}}{22016\mu\pi}, \text{ when } a = 5, b = 4.$$